



Social anxiety in adult males with autism spectrum disorders

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Spain, D., Happé, F., Johnston, P., Campbell, M., Sin, J., Daly, E., Ecker, C., Anson, M., Chaplin, E., Glaser, K., Mendez, A., Lovell, K. and Murphy, D. G. (2016) Social anxiety in adult males with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 32. pp. 13-23. ISSN 1750-9467 doi: <https://doi.org/10.1016/j.rasd.2016.08.002> Available at <http://centaur.reading.ac.uk/71138/>

It is advisable to refer to the publisher's version if you intend to cite from the work.

To link to this article DOI: <http://dx.doi.org/10.1016/j.rasd.2016.08.002>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Social anxiety in adult males with autism spectrum disorders

Author details

Debbie Spain*, MRes, MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Francesca Happé*, PhD, MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Patrick Johnston, PhD, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Malcolm Campbell, PhD, School of Nursing, Midwifery and Social Work, University of Manchester

Jacqueline Sin, PhD, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Eileen Daly, PhD, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Christine Ecker, PhD, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Martin Anson, PhD, South London & Maudsley NHS Foundation Trust, London

Eddie Chaplin, PhD, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London

Karen Glaser, PhD, King's College London

Andreina Mendez, PhD, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London

MRC AIMS Consortium ≠

Karina Lovell§, PhD, School of Nursing, Midwifery and Social Work, University of Manchester

Declan G Murphy§, MD FRCPsych, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology & Neuroscience, King's College London; Sackler Centre for Translational Neurodevelopment, Institute of Psychiatry, Psychology & Neuroscience, King's College London

≠ The MRC AIMS Consortium is a UK collaboration of autism research centres in the UK including the Institute of Psychiatry, Psychology & Neuroscience, London; the Autism Research Centre, University of Cambridge; and the Autism Research Group, University of Oxford. It is funded by the MRC UK and headed by

the Section of Brain Maturation, Institute of Psychiatry, Psychology & Neuroscience. The Consortium members are (in alphabetical order) Bailey AJ, Baron-Cohen S, Bolton PF, Bullmore ET, Carrington S, Chakrabarti B, Daly EM, Deoni SC, Ecker C, Happé F, Henty J, Jezzard P, Johnston P, Jones DK, Lombardo M, Madden A, Mullins D, Murphy C, Murphy DG, Pasco G, Sadek S, Spain D, Steward R, Suckling J, Wheelwright S, and Williams SC.

*Denotes joint first authors

§ Denotes joint senior authors

Keywords

Autism spectrum – Social anxiety – Social phobia – Adults – Self-report questionnaires

Correspondence details

Debbie Spain

Address: MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, De Crespigny Park, PO Box 80, London, SE5 8AF, UK

Telephone: 00 44 207 848 5388

Email: debbie.spain@kcl.ac.uk

Acknowledgments

We would like to thank all the participants who took part in the study. Funding was provided by the Medical Research Council (MRC, UK) and the MRC, UK Autism Imaging Study (AIMS) network (G0400061/69344; DGM, principal investigator). DGM is part supported by the Mortimer D. and Theresa Sackler Foundation, and the National Institute for Health Research (NIHR) Biomedical Research Centre for Mental Health at King's College London, Institute of Psychiatry, Psychology & Neuroscience, and South London and Maudsley National Health Service Foundation Trust. Debbie Spain is funded by an NIHR Clinical Doctoral Research Fellowship (CDRF - 2012 - 03 - 059). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Abstract

Background: Psychiatric conditions, notably anxiety, commonly co-occur with autism spectrum disorders (ASD).

Method: This study investigated self-reported behavioural, cognitive and affective symptoms of social anxiety (SA) in 50 adult males with ASD. Associations between SA, core ASD symptoms and facets of neuropsychological functioning were also examined.

Results: Twenty-six participants (52%) endorsed levels of SA that exceeded the suggested caseness threshold for social anxiety disorder. Categorical and dimensional data analyses indicated that there were no relationships between SA symptoms, present-state or childhood ASD symptom-severity, or measures of socio-emotional processing in this sample.

Conclusions: Study findings suggest that severity of SA is not merely a reflection of ASD symptom-severity. Further research is needed to ascertain the prevalence of SA in adult ASD epidemiological samples, and identify causal and maintaining mechanisms for these co-morbid symptoms.

1. Introduction

Psychiatric disorders are frequently and consistently found to co-occur with autism spectrum disorders (ASD) (e.g. Lever and Geurts, 2016; Russell et al., 2015; Simonoff et al., 2008). High rates and levels of social anxiety, in particular, have been reported in children and adolescents with ASD (e.g. Bellini, 2004; Kuusikko et al., 2008; Melfsen et al., 2006; Russell and Sofronoff, 2005). Data obtained from self- and informant-report instruments suggest that up to 50% of young people with ASD may score above normative levels for social anxiety, although ratings from different informants do not always correlate significantly (Bellini, 2004).

Relatively little is known about social anxiety disorder (SAD) in adults with ASD, despite this being the most common anxiety disorder in the typically developing adult population, with high rates of co-morbid depression, other anxiety disorders, substance use, and increased risk of suicide (NICE, 2013). Cross-sectional studies that have examined general rates of psychiatric co-morbidity in adults with ASD, recruited via community (n=172, Lever and Geurts, 2016) and clinical settings (n=122, Hofvander et al., 2009; n=63, Joshi et al., 2013; n=474, Russell et al., 2015), have estimated that between 12% and 56% of adults meet diagnostic criteria for SAD. Three studies to date, have focused specifically on SAD in adults with ASD. Cath et al. (2008) examined similarities and differences in self-reported SAD, obsessive compulsive disorder (OCD), and affective

symptoms in 12 adults with ASD, compared to matched clinical and non-clinical controls. Participants completed several questionnaires including the Liebowitz Social Anxiety Scale, one of the most widely used self-report social anxiety measures (LSAS: Liebowitz, 1987). Comparable levels of anxiety were found in the SAD, and ASD and SAD groups. Bejerot and colleagues (2014) found that 28% of adults with ASD (n=14 of 50) met the criteria for SAD using the clinician-administered MINI International Neuropsychiatric Interview (M.I.N.I.: Sheehan et al., 1998), as well as the LSAS. Finally, Maddox and White (2015) investigated SAD in three adult samples; individuals with ASD (n=28), individuals with SAD but no ASD (n=26), and non-clinical controls (n=25). Using self-report questionnaires and an objective assessment of anxiety, their findings indicated that 50% of individuals with ASD presented with clinically significant SAD as measured by the Anxiety Disorders Interview Schedule (ADIS-IV: Brown et al., 1994), and the Social Interaction Anxiety Scale (SIAS: Mattick and Clarke, 1998). By contrast, there were no differences between the ASD and ASD+SAD groups on the Brief Fear of Negative Evaluation Scale (Brief FNE: Leary, 1983).

The notion of co-morbid social anxiety in ASD is, however, inherently complex in several respects. First, there is a clear overlap between the symptom profiles of these two disorders (White et al., 2012). ASD is characterised, for example, by qualitative impairments in reciprocal social interaction (WHO, 1992), while hallmark features of SAD also include difficulties with initiating and maintaining interactions and conversations, as well as social avoidance. Second, similar impairments in neuropsychological functioning have been observed in individuals with ASD and those with SAD, such as emotion and face processing deficits (Brunsdon and Happé, 2014; Morrison and Heimberg, 2013; Wong et al., 2012;); again rendering it difficult to demarcate one disorder from the other. Third, both conditions can impair and restrict attainment and independence; symptoms typically affect peer and social relationships, schooling, and employment.

Assessment of SAD in individuals with ASD poses challenges (Kreiser and White, 2014). Individuals with ASD and/or their significant others (e.g. family members) may not spontaneously seek assessment for social avoidance or social evaluative worries, as these characteristics may be attributed to the core disorder. Even when individuals do present to services, impairments in introspection due to theory of mind deficits (Williams and Happé, 2010), or alexithymia (difficulties labelling own emotions, Bird, Press and Richardson, 2011) can render it difficult for them to describe physical and cognitive symptoms of anxiety. Further, while some studies suggest that individuals with ASD are able to self-report psychopathology symptoms (e.g. Berthoz and Hill,

2005; Cadman et al., 2015), commonly used social anxiety measures are yet to be validated for the ASD population. Use of multiple measures that focus on a range of behavioural, cognitive and affective characteristics associated with social anxiety may therefore enhance the screening and assessment process (Kreiser and White, 2014; Maddox and White, 2015; Tyson and Cruess, 2012).

Perhaps as a result of these issues, the relationship between ASD and SAD has seldom been explored. As in typically developing populations, psycho-social factors, including adverse social experiences, cognitive processes such as information and attentional biases, and safety behaviours such as social withdrawal and avoidance, are likely implicated as risk, causal and/or maintaining mechanisms (see Clark, 1999; Morrison and Heimberg, 2013). However, it is also plausible that there are ASD-specific factors that serve to increase vulnerability for, and perpetuate, SAD. For example, it may be that core ASD characteristics, such as deficits in social skills, and/or difficulties with engaging reciprocally in social interaction, contribute to anxiety about social situations (e.g. Bellini, 2004; Tyson and Cruess, 2012; White et al., 2009). Similarly, an intolerance of uncertainty (IoU), or hypo- and hyper-sensory sensitivities, have been found to be associated with anxiety symptoms (Boulter, Freeston, South and Rodgers, 2014; Maisel, Stephenson, South, Rodgers, Freeston and Gaigg, 2016; Wigham, Rodgers, South, McConachie and Freeston, 2015) and these may encourage avoidance of social situations, e.g. because these seem unpredictable or overly stimulating. Additionally, facets of neuropsychological functioning (such as impairments in socio-emotional processing) could be implicated in anxiety development in ASD (White et al., 2009), for example, impairments in the ability to recognise and understand others' thoughts and intentions (Baron-Cohen et al., 2001a), may render social interactions difficult. Finally, poor peer relationships, rejection, and bullying, all of which occur often and repeatedly for young people and adults with ASD (Schroeder et al., 2014), may render this population susceptible to developing social evaluative concerns around difference, inferiority, and vulnerability, as well as encouraging social withdrawal, isolation and avoidance.

In summary, research findings indicate high rates and levels of social anxiety in children and young people with ASD, as measured by self- or informant-based instruments. Few studies have explored the frequency or nature of social anxiety in the adult ASD population, particularly adults who do not have a concurrent intellectual disability, and who are potentially more likely to need to face anxiety-provoking situations in the context of employment or independent living tasks. Also, prevalence estimates have varied widely, which may be due to

differences in study sampling frames and selection criteria, inclusion/exclusion of individuals with heterogeneous ASD presentations, and assessment of SAD using different measures, not all of which rate cognitive, affective and behavioural characteristics associated with social anxiety. Despite the difficulties with assessing and diagnosing SAD in ASD, there is a clear need to better understand if and why these symptoms might co-occur in order to aid early identification of need, and the development of evidence-based treatments.

The aims of the present study were therefore as follows: 1) to explore the frequency and range of self-reported social anxiety symptoms in a sample of adult males with ASD and no intellectual impairment; 2) to examine the relationship between data from multiple self-report social anxiety questionnaires commonly used in clinical/research fields; 3) to investigate the relationship between anxiety symptoms and ASD symptom-severity given that core impairments may be associated with the development of anxiety; and 4) to examine facets of socio-emotional processing in relation to social anxiety. We hypothesised there would be high rates of self-reported social anxiety symptoms, and that there would be associations between social anxiety, ASD symptom-severity, and socio-emotional processing.

2. Methods

2.1 Participants

Participants were recruited from a sample of adult males, living across south-east England, who had previously taken part in the Autism-Imaging case-control Multi-site Study (AIMS: Ecker et al., 2012). The original AIMS sampling frame consisted of 100 males recruited from clinical and non-clinical services (e.g. via ASD non-statutory organisations); 51 of the AIMS participants consented to take part in the present study. Inclusion criteria for the AIMS study were: males aged 18 and over; a clinical-research diagnosis of autism (and no concurrent intellectual impairment) or Asperger's syndrome; and verbal, performance and full scale IQ ≥ 70 . We solely recruited adults who did not have an intellectual impairment, as this could confound results. Individuals were excluded if they had diagnoses of epilepsy, chromosomal or psychotic disorders.

2.2 Materials

Autism spectrum diagnosis: Data pertaining to ASD diagnosis (autism, Asperger syndrome) were obtained from the AIMS dataset. ASD diagnosis was made according to ICD-10 research criteria (WHO, 1992), and confirmed with the ADI-r (Lord et al., 1994). ADI-r scores needed to meet threshold on two of the three domains of ASD

(reciprocal social interaction, communication, and restricted and repetitive patterns of interest and behaviour). Scores could fall below threshold by one point only in one domain (Ecker et al., 2012) given the potential problems with recall when using the ADI-r with adult samples. Present-state assessment of ASD symptomatology was confirmed using the Autism Diagnostic Observation Schedule-generic (ADOS-G: Lord et al., 2000). Clinical and diagnostic assessments were undertaken by psychiatrists or clinical-researchers experienced in working with individuals with ASD; ADI-r and ADOS-G administration were undertaken by reliability-trained clinical-researchers. Participants were also asked to complete the Autism Quotient (AQ; Baron-Cohen et al., 2001b) prior to their initial AIMS appointment.

Social anxiety: Four self-report social anxiety measures were completed. All measures have been validated for non-ASD samples. As normative thresholds have not yet been established for individuals with ASD, we relied on suggested cut-off scores (i.e. denoting clinically significant symptoms) from non-ASD samples, as has been the case for most other studies using self-report measures of psychopathology symptoms in ASD. The primary social anxiety outcome measure for the present study was the Liebowitz Social Anxiety Scale-SR (LSAS-SR: Liebowitz, 1987), as used in two previous studies of SAD in adults with ASD (Bejerot et al., 2014; Cath et al., 2008). The LSAS-SR is a self-report 24 item questionnaire comprising six sub-scales, measuring the extent to which individuals experience fear/anxiety in, and avoid, common social interaction or performance situations, for example “telephoning in public; meeting strangers; and eating in public places”. Items are rated on a four point Likert-scale with a total score of 60 or more suggestive of generalised social anxiety and a maximum score of 144 (Liebowitz, 1987). The LSAS-SR has good psychometric properties in non-ASD samples; internal consistency is high for the total score (α .95), and subscale scores (total fear/anxiety subscale α .91; fear/anxiety in social interaction subscale α .89; fear/anxiety in performance situations subscale α .79; total avoidance subscale α .92; avoidance of social interaction subscale α .89; and avoidance of performance situations subscale α .84) (Baker et al., 2001).

Three further social anxiety questionnaires were administered in order to investigate the range of cognitive, affective and behavioural characteristics associated with social anxiety. The Brief Fear of Negative Evaluation Scale (Brief FNE: Leary, 1983), used in one previous ASD study (Maddox and White, 2015), is a self-report questionnaire rating the strength of belief in cognitions associated with social anxiety, for example “When I am talking to someone, I worry about what they may be thinking of me”. Items are rated on a five point Likert-scale.

A higher score indicates greater social evaluative concerns. There are two principal versions of the Brief FNE: the original version has 12 items (including eight straightforward items and four reversed-scored items), and a more recent eight item version which includes the straightforward items only (Carleton et al., 2011). Participants in the present study completed the 12 item version, although results for the straightforward eight items were calculated and are also reported below. Internal consistency for the 12 item Brief FNE is high in non-ASD samples ($\alpha .90$) (Leary, 1983).

The Social Phobia Scale (SPS: Mattick et al., 1989; Mattick and Clarke, 1998) is a 20 item self-report questionnaire measuring fear associated with being evaluated by others, including items such as “I fear that I may blush when I am with others [and] I am worried people will think my behaviour is odd”. The Social Interaction Anxiety Scale (SIAS: Mattick and Clarke, 1998) is a 20 item self-report questionnaire rating behavioural, affective and cognitive responses during social interaction, for example “I feel tense if I am alone with just one person [and] when mixing socially, I feel uncomfortable”. The SPS and SIAS are typically administered together, although in one recent ASD study, the SIAS was used in isolation (Maddox and White, 2015). Items for both questionnaires are rated on a five point Likert-scale. A higher score suggests greater social anxiety, with a maximum score of 80. The SPS and SIAS clinical cut-off scores suggested by Peters (2000), of >36 and >26 respectively were used. Internal consistency for both measures in (non-ASD) SAD samples is high (SPS $\alpha .89$ and SIAS $\alpha .93$) (Mattick and Clarke, 1998).

General mood and anxiety: Participants also completed a general screening measure of depression and anxiety. The Hospital Anxiety and Depression Scale (HADS: Zigmond and Snaith, 1983) is a self-report 14 item questionnaire measuring anxiety and depression. Items are rated on a four point Likert-scale. A score of eight or more in either subscale indicates caseness, with a maximum score of 21. The English version of the HADS has been used extensively to screen anxiety and depression in non-clinical and clinical samples (but not in ASD samples specifically), with internal consistency ratings of at least $\alpha .76$ (anxiety subscale) and $\alpha .72$ (depression subscale) in non-ASD samples (Bjelland et al., 2002).

IQ: Data pertaining to IQ were obtained from the AIMS dataset. The Wechsler Scale of Intelligence (WASI) was used to estimate verbal, performance, and full scale IQ (Wechsler, 1999).

Socio-emotional processing: We included three measures of socio-emotional processing in the present study, which had also been administered as part of the AIMS study.

The Karolinska Directed Emotional Faces (KDEF: Lundqvist et al., 1998) is a test of emotion recognition comprising 140 natural faces showing happiness, sadness, anger, disgust, fear, surprise, or a neutral expression. Each face is presented with the seven emotion words underneath, and participants are asked to decide which emotion best describes what the person is feeling. Reaction times and the number of correct responses were recorded. The KDEF stimuli have been validated in a non-ASD sample (Goeleven et al., 2008).

The Reading the Mind in the Eyes Task (RMET: Baron-Cohen et al., 2001a): Participants completed an online version of the RMET, comprising 36 photographs of eyes with a choice of four words, from which participants choose the one that best describes what the person in the picture is thinking or feeling. Reaction times and accuracy are recorded. A recent systematic review of the RMET's psychometric properties reported mixed findings in the literature, but the same authors found good internal consistency and test-retest stability in their own validation study (Baron-Cohen et al., 2013).

The Frith-Happé Animations Test (FHA: Castelli et al., 2000; Castelli et al., 2002): This test of mental state attribution ('theory of mind') shows silent animations (39-42 seconds long) of two triangles interacting. Participants were asked to describe what happened in each animation. Data from the 'theory of mind' animations are reported; these are designed to evoke explanations in terms of intentions to deceive, persuade, and so forth. The verbal responses were coded for "intentionality", the degree of mental state attribution (0-5, with absence of mental state language at one pole and elaborate use of mental state language at the other), and "appropriateness" (0-3 with incorrect at one extreme and highly appropriate explanations at the other). Although (good) inter-rater reliability is typically reported in studies using the FHA, the psychometric properties of this experimental measure have not been reported to date.

2.3 Procedure

Participants recruited to the AIMS study completed tasks in the following order: the AQ (and other self-report measures not reported here) were administered via a secure website prior to the initial appointment; the ADI-R was completed by a parent if this had not already been conducted elsewhere; the ADOS-G and WASI were

undertaken at the outset of the testing appointment; and psychometric tasks were completed by participants in a randomised order for counter-balancing purposes (see Wilson et al., 2014 for a comprehensive overview of task administration). The present study used a cross-sectional design. AIMS participants who had consented to be re-contacted for research purposes were asked to complete five self-report questionnaires via a postal survey undertaken between April and August 2010. Attempts to increase the survey response rate included ensuring that the format and readability of questionnaires was clear, along with provision of stamped addressed envelopes, and reimbursement for participation (Edwards et al., 2009). Ethical approvals (REC ref Q0102 / 26) and informed consent were obtained.

2.4 Statistical analyses

Data were analysed using SPSS, version 19 (SPSS Inc.). Continuous variables were assessed with regard to assumptions underlying parametric tests. First, we estimated the reliability (internal consistency) of the social anxiety self-report measures using Cronbach's alpha. We then investigated the frequency and range of social anxiety symptoms for the whole sample. Using correlational analyses, we examined the inter-relation between the different self-report social anxiety questionnaires, and their relationship to ASD symptom-severity as measured by the ADOS-G and ADI-r (clinician-ratings of present-state and childhood ASD symptoms) and the AQ (self-reported ASD symptoms). We also investigated dimensionally the associations between the social anxiety measures and socio-emotional processing. Then, using the LSAS-SR as the primary outcome measure, we divided participants into those scoring above versus below the suggested SAD caseness threshold of 60, and explored whether these groups differed in 1) participant characteristics (age and IQ), 2) ASD symptom-severity, 3) socio-emotional processing, and 4) depression and anxiety scores between the two groups. As no ASD-specific thresholds for SAD on the LSAS-SR have been published, we used current accepted thresholds for the general population to split our ASD participants into those with versus without SAD. Two-sided p values are reported throughout.

3. Results

3.1 Response rate

Fifty-one males consented to complete the questionnaires. Data were excluded for one responding-participant due to missing diagnostic data. There were some missing questionnaire data for a further four participants although we included these individuals in the analyses where possible. We were unable to ascertain reasons for non-

participation in the study, nor were we able to establish the proportion of individuals who were under the care of clinical services. Baseline sample characteristics were compared between individuals who did and did not return questionnaires (see Table 1). There was a significant difference in age between the two groups ($t=2.80$, $df=83.42$, $p=.006$): individuals who did not complete questionnaires were older, on average. There were however, no significant differences in IQ, ADI-r or ADOS-G mean scores ($p>.05$, $d<0.38$) between study participants and those who did not participate.

(Table 1 about here)

3.2 Participant characteristics

Sample characteristics for the 50 male participants are summarised in Table 2. The mean age was 26 years (range 19-42). Full scale IQ for the sample overall was within the average range (IQ=107). Most participants described their ethnicity as 'White European'. A third of participants ($n=17$) were unemployed, 22% ($n=11$) were employed, and 28% ($n=14$) were students. Participants with a diagnosis of Asperger's syndrome did not differ from those with an autism diagnosis on any participant characteristics shown in Table 2, with the exception of higher verbal IQ (mean 113 versus 102), full scale IQ (112 versus 102) and lower ADI-r 'reciprocal social interaction' impairment scores (17 versus 22).

(Table 2 about here)

3.3 Internal consistency of measures

To examine reliability, we estimated Cronbach's alpha for the social anxiety self-report measures used. Internal consistency was high for all the measures: the LSAS-SR (total score α .96; total fear/anxiety subscale α .94; fear/anxiety in social interaction subscale α .92; fear/anxiety in performance situations subscale α .86; total avoidance subscale α .92; avoidance of social interaction subscale α .86; and avoidance of performance situations subscale α .86); the Brief FNE (12 item version α .90; 8 item version α .91); the SPS (α .93); and the SIAS (α .92). All coefficients are comparable to those reported for typically developing samples.

3.4 Analyses for total ASD sample

Frequency of self-reported social anxiety symptoms

Mean scores for each of the four social anxiety questionnaires are outlined in Table 2. Sample scores indicated high levels of self-reported behavioural, cognitive and affective social anxiety symptoms, across all the questionnaires. Using the LSAS-SR as the primary outcome measure, the mean total LSAS-SR score for 46 participants with complete data was 67.3 (s.d. 28.5, range 9-124). The proportion of participants scoring above the LSAS-SR social anxiety threshold was comparable in those diagnosed with autism (65%) and those diagnosed with Asperger's syndrome (52%).

Associations between participant characteristics and social anxiety

There were no significant associations found between the social anxiety measures (total score or subscales of the LSAS-SR, BFNE, SPS or SIAS), and age (all $r < 0.26$; $p > .05$), or verbal, performance, or full scale IQ (all $r < 0.18$; $p > .05$).

Associations between multiple self-report measures of social anxiety

Highly significant correlations were found between the sub-scales of the LSAS-SR (all $r > 0.60$; $p < .005$); and between all other social anxiety measures (all $r > 0.55$; $p < .05$).

Associations between ASD symptom-severity and social anxiety

There were no significant correlations between the social anxiety questionnaires (LSAS, Brief FNE, SPS or SIAS), and domains of the ADI-r (all $r < 0.18$; $p > .05$), or total and subscale scores of the ADOS-G (all $r < 0.10$;

$p > .05$) However, relationships between the AQ and all social anxiety measures were significant (all $r > 0.38$; $p < .04$).

Associations between socio-emotional processing and social anxiety

There were no significant correlations between any of the four social anxiety questionnaires, and the socio-emotional tests (KDEF, RMET and FHA) (all $r < 0.24$; $p > .05$).

3.5 Group differences between ASD participants scoring above versus below the SAD caseness threshold

The sample was divided according to LSAS-SR scores: a threshold score of 60 or more (the suggested threshold for typically developing adult populations) was used to dichotomise the group. Characteristics for individuals with versus without clinical levels of social anxiety on the LSAS-SR are shown in Table 2.

Participant characteristics

Comparing groups scoring above and below the LSAS-SR caseness threshold, there were no significant differences in age ($t = -0.02$, $df = 42$, $p = .986$, $d = -0.07$), verbal IQ ($t = 0.05$, $df = 43$, $p = .961$, $d = 0$), performance IQ ($t = 1.36$, $df = 43$, $p = .180$, $d = 0.44$), or full scale IQ ($t = 0.69$, $df = 43$, $p = .495$, $d = 0.26$).

Associations between ASD symptom-severity and social anxiety

There were no statistically significant differences in mean ASD symptom-severity scores on the ADI-r between groups scoring above versus below the LSAS-SR social anxiety caseness threshold: reciprocal social interaction ($t = 0.38$, $df = 43$, $p = .704$, $d = 0.13$); communication ($t = 0.28$, $df = 43$, $p = .783$, $d = 0.07$); restricted, repetitive and stereotyped behaviours and patterns of interest ($t = 0.96$, $df = 43$, $p = .343$, $d = 0.29$). Differences in the total and subscale scores of the ADOS-G were also not significant: communication ($t = 1.09$, $df = 43$, $p = .283$, $d = 0.33$); reciprocal social interaction ($t = 0.62$, $df = 43$, $p = .542$, $d = 0.18$), total score ($t = 0.80$, $df = 43$, $p = .428$, $d = 0.25$); or stereotyped behaviours and repetitive interests ($t = -0.04$, $df = 43$, $p = .967$, $d = 0$). Similarly, mean scores on the AQ did not differ significantly between groups ($t = -1.58$, $df = 41$, $p = .123$, $d = -0.55$).

Associations between socio-emotional tasks and social anxiety

Performance of participants scoring above versus below the LSAS cut-off for SAD was examined across the three tests of emotion and social cognition. Table 3 shows the scores for these subgroups. There were no

significant differences between mean scores on these measures between the groups: KEDF (RT $t=0.86$, $df=37$, $p=.398$, $d=0.27$; Correct $t=-.09$, $df=37$, $p=.927$, $d=0$); RMET (RT $t=0.12$, $df=39$, $p=.864$, $d=0.05$; Correct $t=0.29$, $df=39$, $p=.773$, $d=0$); and FHA (Intentionality score $t=-1.18$, $df=30$, $p=.247$, $d=-0.5$; Appropriateness score $t=-0.60$, $df=30$, $p=.550$, $d=0$).

(Table 3 about here)

General anxiety and depression scores

Comparing the groups scoring above and below the LSAS-SR caseness threshold, significant differences were found in HADS depression scores ($t=-2.71$, $df=43$, $p=.010$, $d=-0.81$), and anxiety scores ($t=-3.52$, $df=43$, $p=.001$, $d=-1.08$). A significant association was also found between caseness on the LSAS-SR total score and caseness for depression ($\chi^2=6.76$, $df=1$, $p=.009$), and anxiety on the HADS ($\chi^2=7.21$, $df=1$, $p=.007$). Of those scoring in the clinical range for self-reported SAD, 88% ($n=23$) also scored in the clinical range for general anxiety and 54% ($n=14$) in the range for depression.

4. Discussion

While several studies have investigated social anxiety in adults with ASD, there has been limited attention given to potential associations between core ASD characteristics, facets of neuropsychological functioning and social anxiety. The present study investigated social anxiety symptoms, dimensionally and categorically, in a sample of males with ASD. The study also aimed to explore the frequency and range of self-reported social anxiety symptoms in males with ASD, and examine relationships between multiple self-report social anxiety questionnaire measures, and between social anxiety symptoms, ASD symptom-severity and socio-emotional functioning.

First, we found that a significant proportion of participants self-reported social anxiety symptoms across a range of measures. Fifty-two percent of the sample ($n=26$) scored above the suggested caseness threshold on the LSAS-SR. The high self-ratings of social anxiety in our sample are comparable to those reported in younger ASD populations (e.g. Bellini, 2004; Kussikko et al., 2008), a recent adult ASD clinic sample (Joshi et al., 2013), and a combined clinic and community adult sample (who were not reported to be specifically treatment-seeking) (Maddox and White, 2015). These rates are considerably higher than the rates of 7-12% found in epidemiological

studies of typically developing (i.e. non ASD) individuals (NICE, 2013). In relation to previous ASD studies that have employed the LSAS-SR, we found that self-reported social anxiety symptoms were higher in our sample, compared to those reported by Bejerot and colleagues (2014), who found that the mean LSAS-SR score for their sample was 78, and that 28% of participants (n=14) had clinically significant symptoms. Conversely, Cath et al., (2008) reported a mean LSAS-SR score of 107 in their sample of 12 adults.

Are the high levels of social anxiety found in this population simply part and parcel of ASD? Might ratings on social anxiety questionnaires simply be tapping core autism-spectrum features such as impairments in social interaction? The present data suggest not. First, using standardised measures, not all individuals with ASD reached the caseness threshold for social anxiety, despite potential concerns about symptom overlap. Second, measures of social anxiety did not correlate significantly with clinician-rated measures of autism-spectrum symptomatology, nor did ASD symptom-severity on the ADI-r or ADOS differ between subgroups scoring above and below caseness threshold for SAD on the LSAS-SR. This tallies broadly with results from a study of psychiatric co-morbidity in children, in which Simonoff et al. (2008) concluded that autism-severity did not appear to be predictive of co-morbidity (including social phobia). We did find a significant correlation between measures of SAD and the AQ, like Bejerot and colleagues (2014) but unlike that study, we did not find that AQ scores differed in those passing clinical cut-off for SAD on the LSAS-SR. Why the AQ shows a different pattern to the ADOS and ADI-r is uncertain; since the AQ is a self-report measure, common methods variance may be relevant, or SAD may influence self-perceptions of social skills. Indeed, Tonge et al (2016) recently reported elevated AQ scores in (non-ASD) adults with SAD, largely accounted for by items tapping social skills (Tonge et al., 2016). Overall, our results suggest that SAD can co-occur in people with ASD, and be measured beyond the ASD-defining social impairments (although see limitations section below).

Can standard self-report SAD measures be used to screen or aid assessment in individuals with ASD? The present study suggests that they can. Despite possible concerns about difficulties with introspection, the high inter-correlation of the social anxiety measures seems to suggest that social anxiety in ASD may comprise a range of behavioural, cognitive and affective features, as seen in the non-ASD population (NICE, 2013). In addition, the internal consistency of the measures in this ASD sample closely mirrored those reported from typically developing samples.

In the non-ASD adult population, some associations have been found between SAD and facets of social processing, such as deficits in emotion recognition, although findings are not wholly consistent (Morrison and Heimberg, 2013). In the present study, associations between social and emotional tests and self-reported social anxiety were not statistically significant. This may indicate that anxiety is little affected by (current) social cognitive skills, or that the current tasks were not sufficiently sensitive to tap relevant individual differences (e.g. unable to discriminate task-specific compensation from better general adaptation). Further research, using longitudinal study designs, is needed to investigate whether aspects of socio-emotional functioning may contribute to the development or maintenance of social anxiety.

Higher rates and levels of depressive symptoms were endorsed by the group scoring above the SAD caseness threshold. Studies investigating SAD in the typically developing population report similar findings. In a large scale prevalence study, Ohayon and Schatzberg (2010), for example, found that approximately 20% of participants with SAD also met criteria for major depressive disorder. Also, Ghaziuddin and Zafar (2008) and Sterling and colleagues (2008) found that in clinical samples of adults with ASD, depression and anxiety disorders commonly co-occurred. Our findings reiterate the need for future research studies to investigate the potential (inter-dependent) relationships between internalising disorders in the ASD population; for example, do anxiety symptoms contribute to later development of depression, or vice versa?

Several limitations to the study should be noted. First, the sample size, while comparable to previous studies (e.g. Bellini, 2004; Kussikko et al., 2008), is relatively small. To enhance homogeneity of the sample, we only included males, and hence the results may not be applicable to females with ASD. Also, we only included adults with ASD and no intellectual impairments, and so it remains to be seen whether the findings hold true for adults with intellectual disabilities. Second, it was not possible to ascertain the proportion of participants who may have been in receipt of clinical services at the time of study recruitment; nor were data available about socio-economic or employment status, or independent living skills. This may affect the representativeness of the sample, and future research with larger samples well-characterised in these respects, is desirable. Third, only half of those approached returned questionnaires. We cannot be sure whether study participation was affected by participants' experiences of social anxiety. It was noted that study participants and non-responders differed in age: over-representation of younger adults in this study may have skewed the results, and replication is therefore needed in

other age groups. Fourth, despite good psychometric properties of the SAD questionnaires in typically developing samples, they await psychometric validation with the ASD population. Further research is needed to ascertain whether normative thresholds are appropriate for those with ASD, or whether cut-off scores suggestive of clinical caseness should be modified for this group. We also acknowledge that questionnaires were administered in a set order to all participants, and we were therefore unable to explore order effects. Fifth, the study would have been substantially strengthened by use of an objective clinician-administered assessment of SAD, or measures completed by informants (such as parents, carers, or partners). Inclusion of further symptom measures of ASD might also have served to replicate the important finding that SAD symptoms do not appear to be merely a reflection of the severity of ASD symptomatology. Lastly, while correlational analysis provided an estimate of the strength of relationships between measures, causal factors, directions of relationships and confounds could not be assessed in the present study. Future designs, using population-based samples, multiple informants, longitudinal designs and/or intervention trials, could address these issues.

5. Conclusion

This study investigated SAD, using a range of questionnaire measures, in a fairly homogenous sample of adult males with ASD. High rates and levels of social anxiety, general anxiety and low mood were found, corroborating previous findings that internalising disorders are prevalent in this clinical population. Disentangling core ASD characteristics from co-morbid social anxiety symptoms is clearly a complex endeavour for clinicians, researchers, and individuals with ASD (and their significant others), but a failure to consider the co-occurrence of these disorders in routine clinical practice may well leave important needs unassessed and untreated. Further research is now needed to investigate whether bio-psycho-social causal and maintaining factors for social anxiety (in ASD) are similar or distinct to those described for the typically developing population, and to determine what (if any) factors might serve to protect individuals with ASD from developing these co-morbid symptoms.

References

- Baker, S.L., Heinrichs, N., Kim, H-J. & Hofmann, S.G. (2001). The Liebowitz social anxiety scale as a self-report instrument: a preliminary psychometric analysis. *Behaviour Research and Therapy*, 40, 701-715.
- Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J. & Clubley, E. (2001a). The autism-spectrum quotient (AQ). Evidence from Asperger syndrome / high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31, 5-17.
- Baron-Cohen, S., Wheelwright, S., Hill, H., Raste, Y. & Plumb, I. (2001b). The “Reading the Mind in the Eyes” Test Revised Version: A Study with Normal Adults, and Adults with Asperger Syndrome or High Functioning Autism. *Journal of Child Psychology and Psychiatry*, 42(2), 241-251.
- Bejerot, S., Eriksson, J.M. & Mortberg, E. (2014). Social anxiety in adult autism spectrum disorder. *Psychiatry Research*, 220(1), 705-707.
- Bellini, S. (2004). Social Skill Deficits and Anxiety in High Functioning Adolescents with Autism Spectrum Disorders. *Focus on Autism and Other Developmental Disabilities*, 19(2), 78-86.
- Berthoz, S. & Hill, E. (2005). The validity of using self-reports to assess emotion regulation abilities in adults with autism spectrum disorder. *European Psychiatry*, 20, 291-298.
- Bird, G., Press, C. & Richardson, D.C. (2011). The Role of Alexithymia in Reduced Eye-Fixation in Autism Spectrum Conditions. *Journal of Autism and Developmental Disorders*, 41, 1556-1564.
- Bjeland, I., Dahl, A.A., Tangen Haug, T. & Necklemann, D. (2002). The validity of the Hospital Anxiety and Depression Scale: An updated literature review. *Journal of Psychosomatic Research*, 52, 69-77.
- Boulter, C., Freeston, M., South, M. & Rodgers, J. (2014). Intolerance of uncertainty as a framework for understanding anxiety in children and adolescents with autism spectrum disorders. *Journal of Autism and Developmental Disorders*, 44(6), 1391-402.
- Brown, T.A., DiNardo, P.A. & Barlow, D.H. (1994). *Anxiety Disorders Interview Schedule for DSM-IV*. Oxford University Press, New York.
- Brunsdon, V. & Happé, F. (2014). Exploring the ‘fractionation’ of autism at the cognitive level. *Autism*, 18, 17-30.
- Cadman, T., Spain, D., Johnston, P., Russell, A., Mataix-Cols, D., Craig, M. et al. (2015). Obsessive-Compulsive Disorder in Adults with High-Functioning Autism Spectrum Disorder: What Does Self-Report with the OCI-R Tell Us? *Autism Research*, 8, 477-485.

Carleton, R.N., Collimore, K.C., McCabe, R.E. & Antony, M.M. (2011). Addressing revisions to the Brief Fear of Negative Evaluation scale: Measuring fear of negative evaluation across anxiety and mood disorders. *Journal of Anxiety Disorders*, 25, 822-828.

Castelli, F., Frith, C., Happé, F. & Frith, U. (2002). Autism and brain mechanisms for the attribution of mental states to animated shapes. *Brain*, 125: 1839-49.

Castelli, F., Happé, F., Frith, U. & Frith, C. (2000). Movement and mind: a functional imaging study of perception and interpretation of complex intentional movement patterns. *NeuroImage*, 12, 314-325.

Cath, D.C., Ran, N., Smit, J.H., van Balkom, A.J. & Comijs, H.C. (2008). Symptom overlap between autism spectrum disorder, generalized social anxiety disorder and obsessive compulsive disorder in adults: a preliminary case-controlled study. *Psychopathology*, 41, 101-110.

Clark D. M. (1999). A Cognitive Perspective on Social Phobia. In: *International Handbook of Social Anxiety: Concepts, Research and Interventions Relating to the Self and Shyness* (Eds W. R. Crozier and L. E. Alden). 405-430. John Wiley & Sons Ltd., London.

Ecker, C., Suckling, J., Deoni, S.C., Lombardo, M.V., Bullmore, E.T., Baron-Cohen, S. et al. (2012). Brain Anatomy and Its Relationship to Behavior in Adults with Autism Spectrum Disorder. *Archives of General Psychiatry*, 69(2), 195-209.

Edwards, P.J., Roberts, I., Clarke, M.J., DiGiseppi, C., Wentz, R., et al. (2009). *Methods to increase response to postal and electronic questionnaires (Review)*. The Cochrane Collaboration, John Wiley & Sons.

First, M., Spitzer, R., Gibbon, M. & Williams, J. (2002). *Structured Clinical Interview for DSM-IV-TR Axis I Disorders – Patient Edition*.

Ghaziuddin, M. & Zafar, S. (2008). Psychiatric Comorbidity of Adults with Autism Spectrum Disorders. *Clinical Neuropsychiatry*, 5(1), 9-12.

Goeleven, E., Raedt, R.d., Leyman, L., Verschuere, B. (2008). The Karolinska Directed Emotional Faces: A validation study. *Cognition & Emotion*, 22(6), 1094-1118.

Hofvander, B., Delorme, R., Chaste, P., Nyden, A., Wentz, E., Stahlberg, O. et al. (2009). Psychiatric and psychosocial problems in adults with normal-intelligence autism spectrum disorders. *BMC Psychiatry*, 9(35).

Joshi, G., Wozniak, J., Petty, C., Martelon, M.K., Fried, R., Bolfek, A. et al. (2013). Psychiatric comorbidity and functioning in a clinically referred population of adults with autism spectrum disorders: a comparative study. *Journal of Autism and Developmental Disorders*, 43(6), 1314-1325.

Kreiser, N. & White, S.W. (2014). Assessment of Social Anxiety in Children and Adolescents with Autism Spectrum Disorder. *Clinical Psychology Science and Practice*. DOI: 10.1111/cpsp.12057

Kuusikko, S., Pollock-Wurman, R., Jussila, K., Carter, A.S., Mattila, M-L., Ebeling, H. et al. (2008). Social Anxiety in High Functioning Children and Adolescents with Autism and Asperger Syndrome. *Journal of Autism and Developmental Disorders*, 38(9), 1697-1709.

Leary, M.R. (1983). A Brief Version of the Fear of Negative Evaluation Scale. *Personality and Social Psychology Bulletin*, 9(3), 371-375.

Lever, A.G. & Geurts, H.M. (2016) Psychiatric Co-occurring Symptoms and Disorders in Young, Middle-Aged and Older Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 46, 1916-1930.

Liebowitz, M.R. (1987). Social Phobia. *Modern Problems of Pharmacopsychiatry*, 22, 141-173.

Lord, C., Rutter, M., & LeCouteur, A. (1994). Autism Diagnostic Interview-Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 24(5), 659-685.

Lord, C., Risi, S., Leimbrecht, L., Cook, E.H., Leventhal, B.L., DiLavore, P.C. et al. (2000). The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30(3), 205-223.

Lundqvist, D., Flyky, A. & öhman, A. (1998). *The Karolinska Directed Emotional Faces – KDEF, CD ROM*. Department of Clinical Neuroscience, Psychology section, Karolinska Institutet, ISBN 91-630-7164-9.

Maddox B. B. & White S. W. (2015). Comorbid Social Anxiety Disorder in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*. 45(12), 3949-3960.

Maisel, M.E., Stephenson, K.G., South, M., Rodgers, J., Freeston, M.H. & Gaigg, S.B. (2016). Modeling the cognitive mechanisms linking autism symptoms and anxiety in adults. *Journal of Abnormal Psychology*, 125(5), 692-703.

Mattick, R.P. & Clarke, C. (1998). Development and validation of measures of social phobia scrutiny fear and social interaction anxiety. *Behaviour Research and Therapy*, 36, 455-470.

Mattick, R.P., Peters, L. & Clarke., C. (1989). Exposure and Cognitive Restructuring for Social Phobia: A Controlled Study. *Behaviour Therapy*, 20, 3-23.

Melfsen, S., Walita, S. & Warnke, A. (2006). The extent of social anxiety in combination with mental disorders. *European Child and Adolescent Psychiatry*, 15(2), 111-117.

Morrison, A.S. & Heimberg, R.G. (2013). Social Anxiety and Social Anxiety Disorder. *Annual Review of Clinical Psychology*, 9, 249-274.

National Institute for Health and Clinical Excellence (NICE) (2013). *Social Anxiety Disorder: Recognition, Assessment and Treatment*. NICE.

Ohayon, M.M. & Schatzberg, A.F. (2010). Social phobia and depression: prevalence and comorbidity. *Journal of Psychosomatic Research*, 68, 235-243.

Peters, L. (2000). Discriminant validity of the Social Phobia and Anxiety Inventory (SPAI), the Social Phobia Scale (SPS) and the Social Interaction Anxiety Scale (SIAS). *Behaviour Research and Therapy*, 38, 943-950.

Russell, E. & Sofronoff, K. (2005). Anxiety and social worries in children with Asperger syndrome. *Australian and New Zealand Journal of Psychiatry*, 39, 633-638.

Russell, A. J., Murphy, C. M., Wilson, E., Gillan, N., Brown, C., Robertson, D. M. et al., (in press). Short report: the mental health of individuals referred for assessment of Autism Spectrum Disorder (ASD) in adulthood: A clinic report. *Autism*.

Schroeder, J. H., Cappadocia, C. M., Bebko, J. M., Pepler, D. J., & Weiss, J. A. (2014). Shedding Light on a Pervasive Problem: A Review of Research on Bullying Experiences Among Children with Autism Spectrum Disorders. *Journal of Autism and Developmental Disorders*, 44(7), 1520-1534.

Sheehan, D., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E. et al. (1998). The Mini-International Neuropsychiatric Interview (M.I.N.I): The development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *Journal of Clinical Psychiatry*, 59(20), 22-33.

Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T. & Baird, G. (2008). Psychiatric Disorders in Children with Autism Spectrum Disorders: Prevalence, Comorbidity, and Associated Factors in a Population-Derived Sample. *Journal of American Academy of Child and Adolescent Psychiatry*, 47(8), 921-929.

SPSS Inc. (2007). SPSS for Windows statistical software, Release 15. Chicago, Illinois.

Sterling, L., Dawson, G., Estes, A. & Greenson, J. (2009). Characteristics Associated with Presence of Depressive Symptoms in Adults with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 38(6), 1011-1018.

Tonge, N.A., Rodebaugh, T.L., Fernandez, K.C. & Lim, M.H. (2016). Self-reported social skills impairment explains elevated autistic traits in individuals with generalized social anxiety disorder. *Journal of Anxiety Disorders* 38(3), 31-36.

Tyson, K.E. & Cruess, D.G. (2012). Differentiating high-functioning autism and social phobia. *Journal of Autism and Developmental Disorders*, 42(7), 1477-1490.

Vellante, M., Baron-Cohen, S., Melis, M., Marrone, M., Petretto, D.R., & Masala, C. et al. The “Reading the Mind in the Eyes” test: Systematic review of psychometric properties and a validation study in Italy. *Cognitive Neuropsychiatry* (2013) 18(4) 326-354.

Wechsler, D. (1999). *Wechsler Abbreviated Scale of Intelligence (WASI)*. Harcourt Assessment.

White, S.W., Oswald, D., Ollendick, T. & Scahill, L. (2009). Anxiety in children and adolescents with autism spectrum disorders. *Clinical Psychology Review*, 29(3), 216-229.

White, S.W., Bray, B.C. & Ollendick, T.H. (2012). Examining Shared and Unique Aspects of Social Anxiety Disorder and Autism Spectrum Disorder Using Factor Analysis. *Journal of Autism Developmental Disorders*, 42, 874-884.

WHO (1992). *ICD-10*. WHO.

Wigham, S., Rodgers, J., South, M., McConachie, H & Freeston, M. (2015). The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45 (4) 943-952.

Williams, D. & Happé, F. (2010). Representing intentions in self and other: studies of autism and typical development. *Developmental Science*, 13(2), 307-319.

Wilson, C.E., Happé, F., Wheelwright, S., Ecker, C., Lombardo, M.V., Johnston, P et al. (2014). The Neuropsychology of Male Adults with High-Functioning Autism or Asperger Syndrome. *Autism Research*, DOI: 10.1002/aur.1394.

Wong, N., Beidel, D.C., Sarver, D.E. & Sims, V. (2012). Facial Emotion Recognition in Children with High Functioning Autism and Children with Social Phobia. *Child Psychiatry and Human Development*, 43, 775-794.

Zigmond, A.S. & Snaith, R.P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67, 361-370.

Table one – sample characteristics for autism-imaging case-control multi-site study

(AIMS) cohort by response to current study

	Non-responders n=49 mean (s.d.)	Responders n=50 mean (s.d.)
Age	30.3 (8.1)	26.3 (5.8)**
IQ		
<i>Verbal IQ</i>	110 (14.0)	108 (14.9)
<i>Performance IQ</i>	109 (15.8)	105 (15.8)
<i>Full scale IQ</i>	111 (14.6)	108 (14.7)
ADI-R		
<i>Reciprocal social interaction</i>	17.5 (5.0)	19.2 (5.5)
<i>Communication</i>	13.8 (4.3)	14.0 (4.1)
<i>Repetitive behaviours</i>	5.2 (2.4)	4.9 (2.4)
ADOS-G		
<i>Communication</i>	2.9 (1.8)	3.4 (1.7)
<i>Reciprocal social interaction</i>	5.8 (2.9)	6.7 (3.3)
<i>Total ADOS-G score</i>	8.7 (4.1)	10.2 (4.7)
<i>Repetitive behaviours</i>	1.4 (1.5)	0.9 (1.1) ⁺
BAI	12.5 (10.7)	11.0 (10.8)
BDI	12.4 (10.1)	12.1 (10.8)

⁺p=.051

**p≤.01

ADI-R = Autism Diagnostic Interview-Revised; ADOS-G = Autism Diagnostic Observation Schedule; BAI = Beck Anxiety Inventory; BDI = Beck Depression Inventory

Table two – participant characteristics

	Sample n=50	LSAS-SR ≤59 n=19	LSAS-SR ≥ 60 n=26	<i>t</i>	<i>p</i>	<i>d</i>
Age	26.3 (5.8)	26.4 (4.8)	26.8 (6.8)	-0.02	0.986	-0.07
IQ						
<i>Verbal IQ</i>	108 (14.9)	109 (16.3)	109 (14.6)	0.05	0.961	0
<i>Performance IQ</i>	105 (15.8)	109 (17.0)	102 (14.6)	1.36	0.18	0.44
<i>Full scale IQ</i>	107 (14.7)	110 (16.0)	106 (14.5)	0.69	0.495	0.26
ADI-R						
<i>Rec Soc Int</i>	19.2 (5.5)	19.4 (5.4)	18.7 (5.6)	0.38	0.704	0.13
<i>Communication</i>	14.0 (4.1)	14 (4.6)	13.7 (3.7)	0.28	0.783	0.07
<i>Rep behaviours</i>	4.9 (2.4)	5.2 (2.7)	4.5 (2.0)	0.96	0.343	0.29
ADOS-G						
<i>Communication</i>	3.4 (1.7)	3.6 (1.5)	3.1 (1.5)	1.09	0.283	0.33
<i>Rep Soc Int</i>	6.7 (3.3)	6.8 (3.4)	6.2 (3.1)	0.62	0.542	0.18
<i>Total ADOS-G score</i>	10.1 (4.7)	10.4(4.8)	9.3 (4.1)	0.8	0.428	0.25
<i>Rep behaviours</i>	0.9 (1.1)	1.0 (1.2)	1.0 (1.1)	-0.04	0.967	0
AQ	29 (9.2)	26 (10.4)	31 (7.7)	-1.58	0.123	-0.55
LSAS-SR						
<i>Total score</i>	67 (28.5)	40 (12.1)	87 (17.9)	-10.04	.000***	-3.08
<i>Total Fear/Anxiety</i>	35 (15.4)	21 (8.0)	44 (12.1)	-7.1	.000***	-2.24
<i>Total avoidance</i>	33 (15.1)	18 (6.5)	43 (9.7)	-9.72	.000***	-3.03
Brief FNE						
<i>12 items</i>	24 (10.9)	19 (6.8)	28 (11.6)	-3.06	.004**	-0.95
<i>8 items</i>	14 (8.7)	10 (5.7)	18 (8.7)	-3.66	.001***	-1.09
SPS	25 (16.5)	14 (9.6)	33.6 (16.2)	-4.71	.000***	-1.47
SIAS	39 (16.0)	26 (10.4)	49 (12.2)	-6.72	.000***	-2.03
HADS						
<i>Anxiety</i>	10 (5.1)	7.6 (4.3)	12.5 (4.8)	-3.52	.001***	-1.08
<i>Depression</i>	6 (3.8)	4.6 (3.5)	7.3 (3.2)	-2.71	.010**	-0.81

** $p \leq 0.01$

*** $p \leq 0.001$

ADI-R = Autism Diagnostic Interview-Revised; ADOS-G = Autism Diagnostic

Observation Schedule; Brief FNE = Brief Fear of Negative Evaluation Scale;

SPS = Social Phobia Scale; SIAS = Social Interaction Anxiety Inventory;

HADS = Hospital Anxiety and Depression Scale

Table three – Neuropsychological functioning results by caseness on
Liebowitz Social Anxiety Scale (LSAS-SR)

	LSAS-SR ≤59	LSAS-SR ≥ 60
KEDF	(n=16)	(n=23)
<i>RT</i>	3044 (1122)	2793 (707)
<i>Corr</i>	79 (10.1)	79 (10)
RMET	(n=18)	(n=23)
<i>RT</i>	6878 (2429)	6744 (2504)
<i>Corr</i>	22 (4)	22 (7)
FHA	(n=13)	(n=19)
<i>Int</i>	9 (2)	10 (2)
<i>App</i>	3 (2)	3 (2)

KEDF = Karolinska Directed Emotional Faces, Reaction Time and Number Correct;

RMET = Reading the Mind in the Eyes Test, Reaction Time and Number Correct;

FHA = Frith-Happé Animations, Intentionality and Appropriateness